confession!St. Augustine
St. Augustine
coarse-graining

Chapter 9

Transporting densities

what does "anon it lives" refer to?

> Paulina: I'll draw the curtain:
> My lord's almost so far transported that
> He'll think anon it lives.
> W. Shakespeare: *The Winter's Tale*

(P. Cvitanović, R. Artuso, L. Rondoni, and E.A. Spiegel)

In chapters 2, 3, 5 and 6 we learned how to track an individual trajectory, and saw that such a trajectory can be very complicated. In chapter 4 we studied a small neighborhood of a trajectory and learned that such neighborhood can grow exponentially with time, making the concept of tracking an individual trajectory for long times a purely mathematical idealization.

While the trajectory of an individual representative point may be highly convoluted, the density of these points might evolve in a manner that is relatively smooth. The evolution of the density of representative points is for this reason (and other that will emerge in due course) of great interest. So are the behaviors of other properties carried by the evolving swarm of representative points.

We shall now show that the global evolution of the density of representative points is conveniently formulated in terms of evolution operators.

9.1 Measures

> Do I then measure, O my God, and know not what I measure?
> St. Augustine, *The confessions of Saint Augustine*

A fundamental concept in the description of dynamics of a chaotic system is that of *measure*, which we denote by $d\mu(x) = \rho(x)dx$. An intuitive way to define and construct a physically meaningful measure is by a process of *coarse-graining*. Consider a sequence 1, 2, ..., n, ... of increasingly

partition!phase space
phase space!partition
characteristic!function
measure
density



Figure 9.1: (a) First level of partitioning: A coarse partition of $\mathcal{M}$ into regions $\mathcal{M}_0$, $\mathcal{M}_1$, and $\mathcal{M}_2$. (b) $n = 2$ level of partitioning: A refinement of the above partition, with each region $\mathcal{M}_i$ subdivided into $\mathcal{M}_{i0}$, $\mathcal{M}_{i1}$, and $\mathcal{M}_{i2}$.

refined partitions of phase space, figure 9.1, into regions $\mathcal{M}_i$ defined by the characteristic function

$$\chi_i(x) = \begin{cases} 1 & \text{if } x \in \mathcal{M}_i\,, \\ 0 & \text{otherwise}\,. \end{cases} \tag{9.1}$$

A coarse-grained measure is obtained by assigning the "mass", or the fraction of trajectories contained in the ith region $\mathcal{M}_i \subset \mathcal{M}$ at the nth level of partitioning of the phase space:

$$\Delta\mu_i = \int_{\mathcal{M}} d\mu(x)\chi_i(x) = \int_{\mathcal{M}_i} d\mu(x) = \int_{\mathcal{M}_i} dx\,\rho(x)\,. \tag{9.2}$$

The function $\rho(x) = \rho(x,t)$ denotes the *density* of representative points in phase space at time t. This density can be (and in chaotic dynamics, often is) an arbitrarily ugly function, and it may display remarkable singularities; for instance, there may exist directions along which the measure is singular with respect to the Lebesgue measure. As our intent is to sprinkle phase space with a finite number of initial points (repeat an experiment a finite number of times), we shall assume that the measure can be normalized

AH, MAP: define Lebesgue measure

$$\sum_i^{(n)} \Delta\mu_i = 1\,, \tag{9.3}$$

where the sum is over subregions i at the nth level of partitioning. The infinitesimal measure $dx\rho(x)$ can be thought of as an infinitely refined partition limit of $\Delta\mu_i = |\mathcal{M}_i|\rho(x_i)\,,\ x_i \in \mathcal{M}_i$, with normalization

$$\int_{\mathcal{M}} dx\ \rho(x) = 1\,. \tag{9.4}$$

So far, any arbitrary sequence of partitions will do. What are intelligent ways of partitioning phase space? We postpone the answer to chapter 11, after we have developed some intuition about how the dynamics transports densities.

☞ chapter 11

9.2 Perron-Frobenius operator

Jacobian
Dirac delta function
Dirac delta function
delta function!Dirac

Given a density, the question arises as to what it might evolve into with time. Consider a swarm of representative points making up the measure contained in a region $\mathcal{M}_i$ at time $t = 0$. As the flow evolves, this region is carried into $f^t(\mathcal{M}_i)$, as in figure 2.1(b). No trajectory is created or destroyed, so the conservation of representative points requires that

$$\int_{f^t(\mathcal{M}_i)} dx\, \rho(x,t) = \int_{\mathcal{M}_i} dx_0\, \rho(x_0,0)\,.$$

If the flow is invertible and the transformation $x_0 = f^{-t}(x)$ is single-valued, we can transform the integration variable in the expression on the left to

$$\int_{\mathcal{M}_i} dx_0\, \rho(f^t(x_0),t)\left|\det \mathbf{J}^t(x_0)\right|\,.$$

We conclude that the density changes with time as the inverse of the Jacobian (8.19)

$$\rho(x,t) = \frac{\rho(x_0,0)}{|\det \mathbf{J}^t(x_0)|}\,, \qquad x = f^t(x_0)\,, \tag{9.5}$$

which makes sense: the density varies inversely to the infinitesimal volume occupied by the trajectories of the flow.

The manner in which a flow transports densities may be recast into the language of operators, by writing

$$\rho(x,t) = \mathcal{L}^t\rho(x) = \int_{\mathcal{M}} dx_0\, \delta\big(x - f^t(x_0)\big)\, \rho(x_0,0)\,. \tag{9.6}$$

Let us check this formula. Integrating Dirac delta functions is easy: $\int_{\mathcal{M}} dx\, \delta(x) = 1$ if $0 \in \mathcal{M}$, zero otherwise. The integral over a one-dimensional Dirac delta function picks up the Jacobian of its argument evaluated at all of its zeros:

$$\int dx\, \delta(h(x)) = \sum_{\{x:h(x)=0\}} \frac{1}{|h'(x)|}\,, \tag{9.7}$$

and in d dimensions the denominator is replaced by

9.1
page 139

$$\int dx\, \delta(h(x)) = \sum_{\{x:h(x)=0\}} \frac{1}{\left|\det \frac{\partial h(x)}{\partial x}\right|}\,. \tag{9.8}$$

use my transparencies here.

Perron-Frobenius!operator
operator!Perron-Frobenius
transfer!matrix



Figure 9.2: A piecewise-linear skew "Ulam tent" map (9.11) ($\Lambda_0 = 4/3$, $\Lambda_1 = -4$).

Now you can check that (9.6) is just a rewrite of (9.5):

9.2 page 139

$$\begin{aligned} \mathcal{L}^t\rho(x) &= \sum_{x_0=f^{-t}(x)} \frac{\rho(x_0)}{|f^{t\prime}(x_0)|} \qquad \text{(1-dimensional)} \\ &= \sum_{x_0=f^{-t}(x)} \frac{\rho(x_0)}{|\det \mathbf{J}^t(x_0)|} \qquad (d\text{-dimensional})\,. \end{aligned} \tag{9.9}$$

For a deterministic, invertible flow x has only one preimage x_0; allowing for multiple preimages also takes account of noninvertible mappings such as the "stretch & fold" maps of the interval, to be discussed briefly in the next example, and in more detail in sect. 11.3.1.

We shall refer to the kernel of (9.6) as the *Perron-Frobenius operator*:

9.3 page 139

$$\mathcal{L}^t(x,y) = \delta\big(x - f^t(y)\big)\ . \tag{9.10}$$

remember to link with (10.24)

sect. 16.7

remark 15.4

If you do not like the word "kernel" you might prefer to think of $\mathcal{L}^t(x,y)$ as a matrix with indices x, y. The Perron-Frobenius operator assembles the density $\rho(x,t)$ at time t by going back in time to the density $\rho(x_0,0)$ at time $t = 0$.

move this remark to ?: (for nomenclature, see remark 15.4).

in depth: appendix E, p. 801

Example 9.1 Perron-Frobenius operator for a piecewise-linear map: *Consider Assume the expanding 1-d map $f(x)$ of figure 9.2, a piecewise-linear 2–branch map with slopes $\Lambda_0 > 1$ and $\Lambda_1 < -1$:*

9.7 page 141

$$f(x) = \begin{cases} f_0(x) = \Lambda_0 x\,, & x \in \mathcal{M}_0 = [0, 1/\Lambda_0) \\ f_1(x) = \frac{\Lambda_0}{\Lambda_0 - 1}(1-x)\,, & x \in \mathcal{M}_1 = (1/\Lambda_0, 1]\,. \end{cases} \tag{9.11}$$

Both $f(\mathcal{M}_0)$ and $f(\mathcal{M}_1)$ map onto the entire unit interval $\mathcal{M} = [0,1]$. Assume a piecewise constant density

$$\rho(x) = \begin{cases} \rho_0 & \text{if } x \in \mathcal{M}_0 \\ \rho_1 & \text{if } x \in \mathcal{M}_1 \end{cases}\ . \tag{9.12}$$

As can be easily checked using (9.9), the Perron-Frobenius operator acts on this piecewise constant function as a [2×2] "transfer" matrix with matrix elements

stationary!state measure!invariant invariant!measure

9.1 139 9.5 140

duce, explain nsfer" matrix

$$\begin{pmatrix}\rho_0\\ \rho_1\end{pmatrix} \to \mathcal{L}\rho = \begin{pmatrix}\frac{1}{|\Lambda_0|} & \frac{1}{|\Lambda_1|}\\ \frac{1}{|\Lambda_0|} & \frac{1}{|\Lambda_1|}\end{pmatrix}\begin{pmatrix}\rho_0\\ \rho_1\end{pmatrix}, \tag{9.13}$$

stretching both ρ_0 and ρ_1 over the whole unit interval Λ. In this example the density is constant after one iteration, so $\mathcal{L}$ has only a unit eigenvalue $e^{s_0} = 1/|\Lambda_0| + 1/|\Lambda_1| = 1$, with constant density eigenvector $\rho_0 = \rho_1$. The quantities $1/|\Lambda_0|$, $1/|\Lambda_1|$ are, respectively, the fractions of phase space taken up by the $|\mathcal{M}_0|$, $|\mathcal{M}_1|$ intervals. This simple explicit matrix representation of the Perron-Frobenius operator is a consequence of the piecewise linearity of f, and the restriction of the densities ρ to the space of piecewise constant functions. The example gives a flavor of the enterprise upon which we are about to embark in this book, but the full story is much subtler: in general, there will exist no such finite-dimensional representation for the Perron-Frobenius operator. (Continued in example 10.1.)

To a student with a practical bent the example suggests a strategy for constructing evolution operators for smooth maps, as limits of partitions of phase space into regions $\mathcal{M}_i$, with a piecewise-linear approximations f_i to the dynamics in each region, but that would be too naive; much of the physically interesting spectrum would be missed. As we shall see, the choice of function space for ρ is crucial, and the physically motivated choice is a space of smooth functions, rather than the space of piecewise constant functions.

☞ chapter 16

9.3 Invariant measures

A *stationary* or *invariant density* is a density left unchanged by the flow

rethink - where did the Jacobian in (9.5) go?

$$\rho(x,t) = \rho(x,0) = \rho(x)\,. \tag{9.14}$$

Conversely, if such a density exists, the transformation $f^t(x)$ is said to be *measure-preserving*. As we are given deterministic dynamics and our goal is the computation of asymptotic averages of observables, our task is to identify interesting invariant measures for a given $f^t(x)$. Invariant measures remain unaffected by dynamics, so they are fixed points (in the infinite-dimensional function space of ρ densities) of the Perron-Frobenius operator (9.10), with the unit eigenvalue:

repeller measures?

9.3 page 139

$$\mathcal{L}^t\rho(x) = \int_{\mathcal{M}} dy\, \delta(x - f^t(y))\rho(y) = \rho(x). \tag{9.15}$$

⇓PRELIMINARY

For the piecewise linear map example worked out above, in example 10.1, we have already constructed explicitly such eigenfunction, $\rho(y) = $ const., with eigenvalue 1. In general, depending on the choice of $f^t(x)$ and the function space for $\rho(x)$, there may be no, one, or many solutions of the eigenfunction condition (9.15). For instance, a singular measure $d\mu(x) =$

resurect after replacing with the no-escape example the above.

⇑PRELIMINARY

equilibrium!point
confession!C.N. Yang
Yang, C.N.
natural measure
measure!natural
visitation frequency

$\delta(x-x_q)dx$ concentrated on an equilibrium point $x_q = f^t(x_q)$, or any linear combination of such measures, each concentrated on a different equilibrium point, is stationary. There are thus infinitely many stationary measures that can be constructed. Almost all of them are unnatural in the sense that the slightest perturbation will destroy them.

From a physical point of view, there is no way to prepare initial densities which are singular, so it makes sense to concentrate on measures which are limits of transformations experienced by an initial smooth distribution $\rho(x)$ under the action of f, rather than as a limit computed from a single trajectory,

$$\rho_0(x) = \lim_{t\to\infty} \int_{\mathcal{M}} dy\, \delta(x - f^t(y))\rho(y,0)\,, \quad \int_{\mathcal{M}} dy\, \rho(y,0) = 1\,. \tag{9.16}$$

Intuitively, the "natural" measure (or measures) should be the least sensitive to facts of life, such as noise (no matter how weak).

connect to chapter 32: make the robustness under external noise more convincing here

9.3.1 Natural measure

add to refsMeasure.tex

> **Huang:** Chen-Ning, do you think ergodic theory gives us useful insight into the foundation of statistical mechanics?
> **Yang:** I don't think so.
> Kerson Huang, *C.N. Yang interview*

The *natural* or *equilibrium measure* can be defined as the limit

$$\overline{\rho}_{x_0}(y) \;=\; \begin{cases} \lim_{t\to\infty} \frac{1}{t}\int_0^t d\tau\, \delta(y - f^\tau(x_0)) & \text{flows} \\ \lim_{n\to\infty} \frac{1}{n}\sum_{k=0}^{n-1} \delta\big(y - f^k(x_0)\big) & \text{maps}\,, \end{cases} \tag{9.17}$$

9.8 page 141
9.9 page 141

where x_0 is a generic inital point. [1] Staring at an average over infinitely many Dirac deltas is not a prospect we cherish. Generated by the action of f, the natural measure satisfies the stationarity condition (9.15) and is thus invariant by construction. From a computational point of view, the natural measure is the visitation frequency defined by coarse-graining, integrating (9.17) over the $\mathcal{M}_i$ region

$$\Delta\overline{\mu}_i = \lim_{t\to\infty} \frac{t_i}{t}\,, \tag{9.18}$$

where t_i is the accumulated time that a trajectory of total duration t spends in the $\mathcal{M}_i$ region, with the initial point x_0 picked from some smooth density $\rho(x)$.

[1] Driebe: emphasize generic, not periodic: use normal (Borel definition of irrationals) vs. non-normal. Explained in the green book edited by Series

Let $a = a(x)$ be any *observable*. In the mathematical literature $a(x)$ is a function belonging to some function space, for instance the space of integrable functions L^1, that associates to each point in phase space a number or a set of numbers. In physical applications the observable $a(x)$ is necessarily a smooth function. The observable reports on some property of the dynamical system. Several examples will be given in sect. 10.1.

observable
average!space
space!average
functional
time!average
average!time
ergodic!theory
put pointer, refer to the example there
ergodic!average
Birkhoff!ergodic theorem
mixing

The *space average* of the observable a with respect to a measure ρ is given by the d-dimensional integral over the phase space $\mathcal{M}$:

$$\begin{aligned} \langle a \rangle_\rho &= \frac{1}{|\rho_{\mathcal{M}}|} \int_{\mathcal{M}} dx\, \rho(x) a(x) \\ |\rho_{\mathcal{M}}| &= \int_{\mathcal{M}} dx\, \rho(x) = \text{mass in } \mathcal{M}\,. \end{aligned} \tag{9.19}$$

For now we assume that the phase space $\mathcal{M}$ has a finite dimension and a finite volume. By definition, $\langle a \rangle_\rho$ is a function(al) of ρ.

Inserting the right-hand-side of (9.17) into (9.19), we see that the natural measure corresponds to a *time average* of the observable a along a trajectory of the initial point x_0,

$$\overline{a_{x_0}} = \lim_{t\to\infty} \frac{1}{t} \int_0^t d\tau\, a(f^\tau(x_0))\,. \tag{9.20}$$

☞ appendix A

Analysis of the above asymptotic time limit is the central problem of ergodic theory. The Birkhoff ergodic theorem asserts that if a natural measure ρ exists, the limit $\overline{a(x_0)}$ for the time average (9.20) exists for all initial x_0. As we shall not rely on this result in what follows we forgo a proof here. Furthermore, if the dynamical system is *ergodic*, the time average over almost any trajectory tends to the space average

definition of ergodic (Webster?)

$$\lim_{t\to\infty} \frac{1}{t} \int_0^t d\tau\, a(f^\tau(x_0)) = \langle a \rangle \tag{9.21}$$

for "almost all" initial x_0. By "almost all" we mean that the time average is independent of the initial point apart from a set of ρ-measure zero.

explain we drop suffix ρ if ρ is the natural measure

For future reference, we note a further property that is stronger than ergodicity: if the space average of a product of any two variables decorrelates with time,

explain *mixing*

$$\lim_{t\to\infty} \left\langle a(x) b(f^t(x)) \right\rangle = \langle a \rangle \langle b \rangle\ , \tag{9.22}$$

☞ sect. 19.3

the dynamical system is said to be *mixing*.

Henon@Hénon!attractor
deterministic flow
flow!deterministic
stochastic flow
flow!stochastic
evolution!kernel!probabilistic



Figure 9.3: Natural measure (9.18) for the Hénon map (3.12) strange attractor at parameter values $(a, b) = (1.4, 0.3)$. See figure 3.3 for a sketch of the attractor without the natural measure binning. (Courtesy of J.-P. Eckmann)

Example 9.2 The Hénon attractor natural measure: *A numerical calculation of the natural measure (9.18) for the Hénon attractor (3.12) is given by the histogram in figure 9.3. The phase space is partitioned into many equal-size areas $\mathcal{M}_i$, and the coarse grained measure (9.18) is computed by a long-time iteration of the Hénon map, and represented by the height of the column over area $\mathcal{M}_i$. What we see is a typical invariant measure - a complicated, singular function concentrated on a fractal set.*

If an invariant measure is quite singular (for instance a Dirac δ concentrated on a fixed point or a cycle), its existence is most likely of limited physical import. (See however discussion of orbital measures, sect. **??**). No smooth inital density will converge to this measure if the dynamics is unstable. In practice the average (9.17) is problematic and often hard to control, as generic dynamical systems are neither uniformly hyperbolic nor structurally stable: it is not known whether even the simplest model of a strange attractor, the Hénon attractor, is a strange attractor or merely a long stable cycle.

⇓PRELIMINARY
⇑PRELIMINARY

10.1
page 161
Genuflect here to refs. [12.33, 12.34, 12.35]

9.3.2 Determinism vs. stochasticity

While dynamics can lead to very singular ρ's, in any physical setting we cannot do better than to measure it averaged over some region $\mathcal{M}_i$; the coarse-graining is not an approximation but a physical necessity. One is free to think of a measure as a probability density, as long as one keeps in mind the distinction between deterministic and stochastic flows. In deterministic evolution the evolution kernels are not probabilistic; the density of trajectories is transported *deterministically*. What this distinction means will became apparent later: for deterministic flows our trace and determinant formulas will be *exact*, while for quantum and stochastic flows they will only be the leading saddlepoint (stationary phase, steepest descent) approximations.

chapter 15

refer to noise.tex as well
chapter 31

Clearly, while deceptively easy to define, measures spell trouble. The good news is that if you hang on, you will *never need to compute them*, at least not in this book. How so? The evolution operators to which we next turn, and the trace and determinant formulas to which they will lead us, will assign the correct weights to desired averages without recourse to any explicit computation of the coarse-grained measure $\Delta\rho_i$.

9.4 Density evolution for infinitesimal times

continuity equation
semigroup
group!semi-
flow!generator of
generator!of flow

Consider the evolution of a smooth density $\rho(x) = \rho(x,0)$ under an infinitesimal step $\delta\tau$, by expanding the action of $\mathcal{L}^{\delta\tau}$ to linear order in $\delta\tau$:

$$\begin{aligned}
\mathcal{L}^{\delta\tau}\rho(y) &= \int_{\mathcal{M}} dx\, \delta\Big(y - f^{\delta\tau}(x)\Big)\, \rho(x) \\
&= \int_{\mathcal{M}} dx\, \delta(y - x - \delta\tau v(x))\, \rho(x) \\
&= \frac{\rho(y - \delta\tau v(y))}{\left|\det\left(1 + \delta\tau \frac{\partial v(y)}{\partial x}\right)\right|} = \frac{\rho(y) - \delta\tau \sum_{i=1}^{d} v_i(y)\partial_i \rho(y)}{1 + \delta\tau \sum_{i=1}^{d} \partial_i v_i(y)} \\
\rho(x,\delta\tau) &= \rho(x,0) - \delta\tau \sum_{i=1}^{d} \frac{\partial}{\partial x_i}(v_i(x)\rho(x,0))\,. \qquad (9.23)
\end{aligned}$$

Here we have used the infinitesimal form of the flow (2.5), the Dirac delta Jacobian (9.9), and the $ln\, det = tr\, ln$ relation. Moving $\rho(y,0)$ to the left hand side and dividing by $\delta\tau$, we discover that the rate of the deformation of ρ under the infinitesimal action of the Perron-Frobenius operator is nothing but the *continuity equation* for the density:

4.1
page 73

$$\partial_t \rho + \partial \cdot (\rho v) = 0\,. \qquad (9.24)$$

The family of Perron-Frobenius operators operators $\{\mathcal{L}^t\}_{t\in\mathbb{R}_+}$ forms a semigroup parametrized by time

(a) $\mathcal{L}^0 = I$

(b) $\mathcal{L}^t\mathcal{L}^{t'} = \mathcal{L}^{t+t'} \qquad t, t' \geq 0 \qquad$ (semigroup property) .

The *generator* of the semigroup (that is, the generator of a time translation by an infinitesimal step $\delta\tau$) is determined by (9.23):

$$\mathcal{A}\rho(x) = +\lim_{\delta\tau\to 0^+} \frac{1}{\delta\tau}\left(\mathcal{L}^{\delta\tau} - I\right)\rho(x) = -\partial_i(v_i(x)\rho(x))\,. \qquad (9.25)$$

looks like Hamiltonian, but it is not

If the flow is finite-dimensional and invertible, $\mathcal{A}$ is a generator of a full-fledged group. The left hand side is the definition of time derivative, so the evolution equation for $\rho(x)$ is

$$\left(\frac{\partial}{\partial t} - \mathcal{A}\right)\rho(x) = 0\,. \qquad (9.26)$$

appendix E.2

The finite time Perron-Frobenius operator (9.10) can be formally expressed by exponentiating the time evolution generator $\mathcal{A}$ as

$$\mathcal{L}^t = e^{t\mathcal{A}}\,. \qquad (9.27)$$

operator!semigroup!bounded
semigroup!operator
Laplace!transform
resolvent!operator
operator!resolvent

9.10
page 141

⇓PRELIMINARY
??
page ??
⇑PRELIMINARY

The generator $\mathcal{A}$ is reminiscent of the generator of translations. Indeed, for a constant velocity field dynamical evolution is nothing but a translation by (time $\times$ velocity):

$$e^{-tv\frac{\partial}{\partial x}}a(x) = a(x - tv)\,. \tag{9.28}$$

As we will not need to implement a computational formula for general $e^{t\mathcal{A}}$ in what follows, we relegate making sense of such operators to appendix E.2.

☞ appendix E.2

9.4.1 Resolvent of $\mathcal{L}$

Here we limit ourselves to a brief remark about the notion of the "spectrum" of a linear operator.

The Perron-Frobenius operator $\mathcal{L}$ acts multiplicatively in time, so it is reasonable to suppose that there exist constants $M > 0$, $\beta \geq 0$ such that $||\mathcal{L}^t|| \leq Me^{t\beta}$ for all $t \geq 0$. What does that mean? The operator norm is defined in the same spirit in which one defines matrix norms (see appendix M.2): We are assuming that no value of $\mathcal{L}^t\rho(x)$ grows faster than exponentially for any choice of function $\rho(x)$, so that the fastest possible growth can be bounded by $e^{t\beta}$, a reasonable expectation in the light of the simplest example studied so far, the exact escape rate (10.20). If that is so, multiplying $\mathcal{L}^t$ by $e^{-t\beta}$ we construct a new operator $e^{-t\beta}\mathcal{L}^t = e^{t(\mathcal{A}-\beta)}$ which decays exponentially for large t, $||e^{t(\mathcal{A}-\beta)}|| \leq M$. We say that $e^{-t\beta}\mathcal{L}^t$ is an element of a *bounded* semigroup with generator $\mathcal{A} - \beta I$. Given this bound, it follows by the Laplace transform

☞ appendix M.2

EAS: need to define the norm of this operator

$$\int_0^\infty dt\, e^{-st}\mathcal{L}^t = \frac{1}{s-\mathcal{A}}\,, \qquad \mathrm{Re}\, s > \beta\,, \tag{9.29}$$

$\mathcal{A}$ sign wrong here

that the *resolvent* operator $(s-\mathcal{A})^{-1}$ is bounded ("resolvent" = able to cause separation into constituents)

☞ sect. M.2

maybe also as Fourier transform of (9.26), a "Green's" function

$$\left|\left|\frac{1}{s-\mathcal{A}}\right|\right| \leq \int_0^\infty dt\, e^{-st}Me^{t\beta} = \frac{M}{s-\beta}\,.$$

If one is interested in the spectrum of $\mathcal{L}$, as we will be, the resolvent operator is a natural object to study. The main lesson of this brief aside is that for continuous time flows, the Laplace transform is the tool that brings down the generator in (9.27) into the resolvent form (9.29) and enables us to study its spectrum.

in depth:
appendix E.2, p. 803

9.5 Liouville operator

Poisson!bracket
flow!incompressible
conservation!phase space volume
Liouville!theorem
density!phase space
phase space!density
continuity!equation

A case of special interest is the Hamiltonian or symplectic flow defined by the Hamilton's equations of motion (5.1). A reader versed in quantum mechanics will have observed by now that with replacement $\mathcal{A} \to -\frac{i}{\hbar}\hat{H}$, where $\hat{H}$ is the quantum Hamiltonian operator, (9.26) looks rather like the time dependent Schrödinger equation, so this is probably the right moment to figure out what all this means in the case of Hamiltonian flows.

The Hamilton's evolution equations (5.1) for any time-independent quantity $Q = Q(q,p)$ are given by

$$\frac{dQ}{dt} = \frac{\partial Q}{\partial q_i}\frac{dq_i}{dt} + \frac{\partial Q}{\partial p_i}\frac{dp_i}{dt} = \frac{\partial H}{\partial p_i}\frac{\partial Q}{\partial q_i} - \frac{\partial Q}{\partial p_i}\frac{\partial H}{\partial q_i}\,. \tag{9.30}$$

As equations with this structure arise frequently for symplectic flows, it is convenient to introduce a notation for them, the *Poisson bracket*

$$\{A,B\} = \frac{\partial A}{\partial p_i}\frac{\partial B}{\partial q_i} - \frac{\partial A}{\partial q_i}\frac{\partial B}{\partial p_i}\,. \tag{9.31}$$

Gaspard has $\{A,B\}$ defined the other way around; use $\{,\}$

In terms of Poisson brackets the time evolution equation (9.30) takes the compact form

$$\frac{dQ}{dt} = \{H,Q\}\,. \tag{9.32}$$

The full phase space flow velocity is $\dot{x} = (\dot{q},\dot{p})$, where the dot signifies time derivative for fixed initial point.

The discussion of sect. 9.4 applies to any deterministic flow. If the density itself is a material invariant, combining (??) and (9.24) we conclude that $\partial_i v_i = 0$ and $\mathbf{J}^t(x_0) = 1$. An example of such incompressible flow is the Hamiltonian flow of sect. 5.2. For incompressible flows the continuity equation (9.24) becomes a statement of conservation of the phase space volume (see sect. 5.2), or the *Liouville theorem*

⇓PRIVATE
⇑PRIVATE

$$\partial_t \rho + v_i \partial_i \rho = 0\,. \tag{9.33}$$

Hamilton's equations (5.1) imply that the flow is incompressible, $\partial_i v_i = 0$, so for Hamiltonian flows the equation for ρ reduces to the *continuity equation* for the phase-space density:

☞ appendix E

confused - conflicts with (9.24) and (9.33)

$$\partial_t \rho + \partial_i(\rho v_i) = 0\,. \tag{9.34}$$

Consider the evolution of the phase space density ρ of an ensemble of noninteracting particles; the particles are conserved, so

reference is Goldstein

Liouville!equation
Poisson!bracket
symplectic!transformation
Liouville!operator
operator!Liouville
recheck sign!

$$\frac{d}{dt}\rho(q,p,t) = \left(\frac{\partial}{\partial t} + \dot{q}_i\frac{\partial}{\partial q_i} + \dot{p}_i\frac{\partial}{\partial p_i}\right)\rho(q,p,t) = 0\,.$$

Inserting Hamilton's equations (5.1) we obtain the *Liouville equation*, a special case of (9.26):

$$\frac{\partial}{\partial t}\rho(q,p,t) = -\mathcal{A}\rho(q,p,t) = \{H,\rho(q,p,t)\}\,, \tag{9.35}$$

rescue the Poisson brackets from appendix D

where $\{\ ,\ \}$ is the Poisson bracket (9.31). The generator of the flow (9.25) is now the generator of infinitesimal symplectic transformations,

$$\mathcal{A} = \dot{q}_i\frac{\partial}{\partial q_i} + \dot{p}_i\frac{\partial}{\partial p_i} = \frac{\partial H}{\partial p_i}\frac{\partial}{\partial q_i} - \frac{\partial H}{\partial q_i}\frac{\partial}{\partial p_i}\,. \tag{9.36}$$

photoshop

For separable Hamiltonians of form $H = p^2/2m + V(q)$, the equations of motion are

$$\dot{q}_i = \frac{p_i}{m}\,, \qquad \dot{p}_i = -\frac{\partial V(q)}{\partial q_i}\,. \tag{9.37}$$

move to appendix F

and the action of the generator

$$\mathcal{A} = -\frac{p_i}{m}\frac{\partial}{\partial q_i} + \partial_i V(q)\frac{\partial}{\partial p_i}\,. \tag{9.38}$$

9.11
page 141

Dorfman notation is $L = -\mathcal{A}$

can be interpreted as a translation (9.28) in configuration space, followed by acceleration by force $\partial V(q)$ in the momentum space.

This special case of the time evolution generator (9.25) for the case of symplectic flows is called the *Liouville operator*. You might have encountered it in statistical mechanics, while discussing what ergodicity means for 10^{23} hard balls. Here its action will be very tangible; we shall apply the evolution operator to systems as small as 1 or 2 hard balls and to our surprise learn that that suffices to alredy get a bit of a grip on foundations of the classical nonequilibrium statistical mechanics.

in depth:
sect. E.2, p. 803

Commentary

Remark 9.1 Ergodic theory. An overview of ergodic theory is outside the scope of this book: the interested reader may find it useful to consult [9.1]. The existence of time average (9.20) is the basic result of ergodic theory, known as the Birkhoff theorem, see for example refs. [9.1, 1.15], or the statement of theorem 7.3.1 in ref. [9.6]. The natural measure (9.18) (more carefully defined than in the above sketch) is often referred to as the SRB or Sinai-Ruelle-Bowen measure [1.18, 1.17, 1.20].

Remark 9.2 Bounded semigroup. For a discussion of the bounded semigroups of page 134 see, for example, Marsden and Hughes [9.2].

Poisson!bracket

Remark 9.3 The sign convention of the Poisson bracket. The Poisson bracket is antisymmetric in its arguments and there is a freedom to define it with either sign convention. When such freedom exists, it is certain that both conventions are in use and this is no exception. In some texts [1.4, 9.3] you will see the right hand side of (9.31) defined as $\{B, A\}$ so that (9.32) is $\frac{dQ}{dt} = \{Q, H\}$. Other equally reputable texts [37.27] employ the convention used here. Landau and Lifshitz [9.4] denote a Poisson bracket by $[A, B]$, notation that we reserve here for the quantum-mechanical commutator. As long as one is consistent, there should be no problem.

Résumé

In physically realistic settings the initial state of a system can be specified only to a finite precision. If the dynamics is chaotic, it is not possible to calculate accurately the long time trajectory of a given initial point. Depending on the desired precision, and given a deterministic law of evolution, the state of the system can then be tracked for a finite time.

The study of long-time dynamics thus requires trading in the evolution of a single phase space point for the evolution of a *measure*, or the *density* of representative points in phase space, acted upon by an *evolution operator*. Essentially this means trading in nonlinear dynamical equations on finite dimensional spaces $x = (x_1, x_2 \cdots x_d)$ for linear equations on infinite dimensional vector spaces of density functions $\rho(x)$. The most physical of stationary measures is the natural measure, a measure robust under perturbations by weak noise.

Reformulated this way, classical dynamics takes on a distinctly quantum-mechanical flavor. If the Lyapunov time (1.1), the time after which the notion of an individual deterministic trajectory loses meaning, is much shorter than the observation time, the "sharp" observables are those dual to time, the eigenvalues of evolution operators. This is very much the same situation as in quantum mechanics; as atomic time scales are so short, what is measured is the energy, the quantum-mechanical observable dual to the time. For long times the dynamics is described in terms of stationary measures, that is, fixed points of certain evolution operators. Both in classical and quantum mechanics one has a choice of implementing dynamical evolution on densities ("Schrödinger picture", sect. 9.4) or on observables ("Heisenberg picture", sect. 10.2 and chapter 14). By "Schrödinger picture" we mean computer eigenvalues...., and then expectation values by extreme sandwiching the observables between the eigen.....

⇓PRELIMINARY

⇑PRELIMINARY

In what follows we shall find the second formulation more convenient, but the alternative is worth keeping in mind when posing and solving invariant density problems. However, as classical evolution operators are not

unitary, their eigenstates can be quite singular and difficult to work with. In what follows we shall learn how to avoid dealing with these eigenstates altogether. As a matter of fact, what follows will be a labor of radical deconstruction; after having argued so strenuously here that only smooth measures are "natural", we shall merrily proceed to erect the whole edifice of our theory on periodic orbits, that is, objects that are δ-functions in phase space. The trick is that each comes with an interval, its neighborhood – cycle points only serve to pin these intervals, just as the millimeter marks on a measuring rod partition continuum into intervals.

References

need more realistic set of refs on ergodic theory

[9.1] Ya.G. Sinai, *Topics in Ergodic Theory* (Princeton Univ. Press, Princeton, New Jersey 1994). [2]

[9.2] J.E. Marsden and T.J.R. Hughes, *Mathematical Foundations of Elasticity* (Prentice-Hall, Englewood Cliffs, New Jersey 1983)

[9.3] H. Goldstein, *Classical Mechanics* (Addison-Wesley, Reading, 1980).

[9.4] L.D. Landau and E.M. Lifshitz, *Mechanics* (Pergamon, London, 1959).

[9.5] P. Cvitanović, C.P. Dettmann, R. Mainieri and G. Vattay, *Trace formulas for stochastic evolution operators: Weak noise perturbation theory*, *J. Stat. Phys.* **93**, 981 (1998); `chao-dyn/9807034`.

[9.6] A. Lasota and M.C. Mackey, *Chaos, Fractals and Noise* (Springer, New York 1994).

[9.7] We thank Laurette Tuckerman for providing us with the pseudospectral code.

[9.8] We have used the NAG library integration routine D02BHF. Our code and numerical results are available on `ChaosBook.org/extras`.

[2] Mason: Sinai has a more elementary book on this