Dynamical systems approach to 1-d spatiotemporal chaos – A cyclist's view

A Thesis
Presented to
The Academic Faculty

by

Yueheng Lan

In Partial Fulfillment
of the Requirements for the Degree
Doctor of Philosophy

School of Physics
Georgia Institute of Technology
target submission date: May 18, 2004

Dynamical systems approach to 1-d spatiotemporal chaos –
A cyclist's view

Approved by:

Professor Chair, Committee Chair	Professor 2nd Reference (Another Department)
Professor Predrag Cvitanović, Adviser	Professor 3rd
Professor 1st	Professor 4th

Date Approved _____

TABLE OF CONTENTS

LIST OF TABLES . . . . . **vii**

LIST OF FIGURES . . . . . **viii**

SUMMARY . . . . . **xi**

I INTRODUCTION **1**

1.1 From dynamical systems to periodic orbit theory 1

1.1.1 Historical perspective 1

1.1.2 Different approaches to dynamical systems 2

1.2 Periodic orbits: theory and numerics 3

1.3 Turbulence:brief notes 5

1.3.1 Historical perspective 5

1.3.2 Main lines of approach 6

1.3.3 Spatiotemporal chaos 7

1.4 Periodic orbit theory on spatiotemporal chaos 8

II PERIODIC ORBIT THEORY **10**

2.1 Dynamical systems 10

2.2 Physical averages in a dynamical system 12

2.2.1 Space, time averages and ergodicity 12

2.2.2 Evolution operators and invariant measures 13

2.3 Trace formula, spectral determinant and dynamical zeta function 14

2.3.1 Trace formula 14

2.3.2 Geometrical interpretation 17

2.3.3 Correlation functions 18

2.4 Cycle expansions 19

2.4.1 Cycle formula for dynamical averages 21

2.5 Summary 22

III PERIODIC ORBITS AND HOW TO FIND THEM **24**

3.1 Review of known techniques for locating UPOs 24

3.1.1 Inverse iteration and anti-integral limit method 24

3.1.2 Phase-space partition method and cyles from long time series . . . 26
3.1.3 Shooting method . . . 26
3.1.4 Variational and relaxation approach . . . 27
3.2 The Newton descent method in loop space . . . 29
3.2.1 A variational equation for the loop evolution . . . 29
3.2.2 Marginal directions and accumulation of loop points . . . 32
3.3 Extensions of Newton descent . . . 33
3.4 Implementation of Newton descent . . . 35
3.4.1 Numerical implementation . . . 35
3.4.2 Initialization and convergence . . . 37
3.4.3 Symmetry considerations . . . 38
3.5 Applications . . . 39
3.5.1 Hénon-Heiles system and restricted three-body problem . . . 39
3.5.2 Periodic orbits of Kuramoto-Sivashinsky system . . . 40
3.6 Summary . . . 42
IV COMPLEX GINZBURG-LANDAU EQUATION . . . 44
4.1 Review . . . 44
4.1.1 Introduction . . . 44
4.1.2 Symmetries and Bekki-Nozaki solutions . . . 45
4.1.3 Stokes solutions and their stability . . . 46
4.1.4 Low-dimensional behavior of the CGLe . . . 47
4.1.5 Phase chaos versus defect chaos . . . 48
4.2 Coherent structures and MAWs . . . 49
4.3 Reformulation of the problem . . . 50
4.4 4-D dynamical system and the existence of periodic solutions . . . 53
4.4.1 Case I . . . 55
4.4.2 Case II . . . 58
4.5 Analytic form of periodic solutions, stability analysis and numerical tests . 60
4.5.1 Case I . . . 61
4.5.2 Case II . . . 63
4.6 Summary . . . 66

V 1 – D KURAMOTO-SIVASHINSKY EQUATION AND ITS STEADY SOLUTIONS 68

5.1 Derivation 68

5.2 Basic Properties 70

5.2.1 Symmetry and Fourier modes 70

5.2.2 Origin is the global attractor for small L 71

5.2.3 Inertial manifold 72

5.3 Steady solutions of the KSe 72

5.3.1 Heteroclinic and homoclinic Connections 74

5.3.2 Periodic Solutions of the steady equation 75

5.3.3 Phase space structure at small c 77

5.3.4 Variational approach 79

5.3.5 Other bounded solutions 80

5.4 Summary 81

VI SPATIOTEMPORAL DYNAMICS OF THE 1-D KURAMOTO-SIVASHINSKY EQUATION 82

6.1 Review 82

6.1.1 Bifurcation sequence 82

6.1.2 Spatiotemporal chaos and recurrence patterns 83

6.2 Dynamics of KSe at $L = 38.5\,, \nu = 1$ 84

6.2.1 Qualitative dynamics and Fourier modes 84

6.2.2 Steady solutions and the dynamics 85

6.2.3 Poincaré sections and intrinsic coordinates 86

6.2.4 Return maps and periodic orbits 88

6.3 Summary 90

VII SUMMARY 93

APPENDIX A — REMARKS ON THE JACOBIANS FOR THE GENERAL FLOW 95

APPENDIX B — COMPARISON OF NEWTON DESCENT WITH VARIATIONAL PRINCIPLE IN HAMILTONIAN SYSTEM 97

APPENDIX C — THEOREMS USED IN THE PROOFS 101

REFERENCES **104**

INDEX **121**

LIST OF TABLES

LIST OF FIGURES

1 The Ulam map $f(x) = 4x(1-x)$ on the unit interval. (a) The graph of the map with its symbolic regions denoted by 0 and 1. (b) The invariant measure: analytic result is depicted by the solid line; numerical result by the dots. 11

2 (a) An annulus $L(\tau)$ swept by the Newton descent flow $d\tilde{x}/d\tau$, connecting smoothly the initial loop $L(0)$ to the periodic orbit $p = L(\infty)$. (b) In general the loop velocity field $\tilde{v}(\tilde{x})$ does not coincide with $\lambda v(\tilde{x})$; for a periodic orbit p, it does so at every $x \in p$. 31

3 The Hénon-Heiles system in a chaotic region: (a) An initial loop $L(0)$, and (b) the unstable periodic orbit p of period $T = 13.1947$ reached by the Newton descent (60). (c) The exponential decrease of the cost function, $\ln(F^2) \approx -2.0502\,\tau + 6.0214$. 39

4 (a) An initial loop $L(0)$, and (b) the unstable periodic orbit p of period $T_p = 2.7365$ reached by the Newton descent (60), the restricted three body problem (66) in the chaotic regime, $\mu = 0.04$. 40

5 The Kuramoto-Sivashinsky system in a spatio-temporally turbulent regime (viscosity parameter $\nu = 0.015$, $d = 32$ Fourier modes truncation). (a) An initial guess L_1, and (b) the periodic orbit p_1 of period $T_1 = 0.744892$ reached by the Newton descent. (c) Another initia

6 Level plot of the space-time evolution $u(x,t)$ for the two spatio-temporally periodic solutions of figure 5: (a) the evolution of p_1, with start of a repeat after the cycle period $T_1 = 0.744892$, and (b) one full period $T_2 = 1.184668$ in the evolution of p_2. 42

7 A typical evolution of the CGLe (70) in the chaotic regime in a periodic box with size $L = 100$. $\alpha = 1.5\,, \beta = -1.4\,, 128$ Fourier modes used. (a) The space-time gray scale plot; (b) The spatial profile of $|A|$ at $t = 450$. 45

8 Marginal stability curve (MS) and Eckhaus instability curve (E) defining regions where plane waves solutions exist in the CGLe (inside (MS)), and are stable when $1 + \alpha\beta > 0$ (inside (E)). Case I corresponds to the wave of maximal possible wavenumber and Case II to t

9 Phase diagram of the CGLe. The line L_{BF} is the locus of the Benjamin-Feir instability. L_1, L_2 and L_3 intersect at the point C^*. Below L_1 and L_3, only defect chaos is observed. The region enclosed by L_2, L_3 and L_{BF} is a bichaotic region (marked BC). Note that L_2 cro

10 Left: wavenumber distribution of stationary MAWs in the (α, β) plane. In (BFS), MAWs exist if $q^2 > q_M^2$. In (M1), MAWs exist $\forall q$. In (M2), MAWs exist if $q^2 < q_M^2$. Right: regions of existence of MAWs in the (q, μ) plane in the Benjamin-Feir-Newell stable regime ((BF

11 Spatial profile of the amplitude $R(x)$ from (117) at $\mu = 0.07264$, with $R_{max} = 0.075$. 62

12 Spatial profiles of the amplitude R for $\mu = 1, \alpha = 1.5, \beta = -1.2, L = 8.958$ from numerical simulation (dots) and the approximate solution (119) (solid line). The agreement is good, with the discrepancy mainly due to the long relaxation time close to the bifurcation point

13 The stability eigenvalue of $u(x,t) \equiv 0$ versus the wavenumber kq. 71

14 Phase diagram of (148) to the lowest order. 78

15 A non-symmetric steady solutions of (135),$L = 38.5\,, \nu = 1\,$. (a) u, u_x projection in the 3-d phase space of (139); (b) The profile of the same orbit on the interval $[0, L]$; 79

16 The first antisymmetric steady solutions of (135),$L = 38.5\,, \nu = 1\,$. (a) u, u_x projection in the 3-d phase space of (139); (b) The profile of the same orbit on the interval $[0, L]$; 79

17 The second antisymmetric steady solutions of (135),$L = 38.5\,, \nu = 1\,$. (a) u, u_x projection in the 3-d phase space of (139); (b) The profile of the same orbit on the interval $[0, L]$; 80

18 The third antisymmetric steady solution of (135),$L = 38.5\,, \nu = 1\,$. (a) u, u_x projection in the 3-d phase space of (139); (b) The profile of the same orbit on the interval $[0, L]$; 80

19 The fourth antisymmetric steady solution of (135),$L = 38.5\,, \nu = 1\,$. (a) u, u_x projection in the 3-d phase space of (139); (b) The profile of the same orbit on the interval $[0, L]$; 81

20 The fifth antisymmetric steady solutions of (135),$L = 38.5\,, \nu = 1\,$. (a) u, u_x projection in the 3-d phase space of (139); (b) The profile of the same orbit on the interval $[0, L]$; 81

21 A sketch of the bifurcation diagram of the KSe in the antisymmetric solution space with periodic boundary conditions. The system size L is the bifurcation parameter. 83

22 Large scale space-time evolution of the KSe in the full solution space with periodic boundary condition, 512 Fourier modes truncation, $L = 400\,, \nu = 1$. 84

23 (a) a_1, a_2 projection of a typical phase orbit of (135); (b) a_3, a_4 projection of the same orbit; $L = 38.5\,, \nu = 1\,,$ 16 Fourier modes truncation. 85

24 The spatial structure of $u(x,t)$ at a point on S_R, $L = 38.5\,, \nu = 1\,,$ 16 Fourier modes truncation: (a) $u\,, u_x$ representation of the state; (b) Spatial profile of the same state. 86

25 The spatial structure of $u(x,t)$ at a point on S_C, $L = 38.5\,, \nu = 1\,,$ 16 Fourier modes truncation: (a) $u\,, u_x$ representation of the state; (b) Spatial profile of the same state. 86

26 The spatial structure of $u(x,t)$ at a point on the transition line from S_R to S_C, $L = 38.5\,, \nu = 1\,,$ 16 Fourier modes truncation: (a) $u\,, u_x$ representation of the state; (b) Spatial profile of the same state. 87

27 The shortest periodic orbit p_C in S_C, $T = 25.6356\,, L = 38.5\,, \nu = 1\,$. (a) $a_1\,, a_2$ Fourier modes projection of the orbit; (b) $a_3\,, a_4$ projection of the same orbit. 87

28 The first shortest periodic orbit p_{R1} in S_R, $T = 20.0229\,, L = 38.5\,, \nu = 1\,$. (a) $a_1\,, a_2$ Fourier modes projection of the orbit; (b) $a_3\,, a_4$ projection of the same orbit. 88

29 The second shortest periodic orbit p_{R2} in S_R, $T = 12.0798\,, L = 38.5\,, \nu = 1\,$. (a) $a_1\,, a_2$ Fourier modes projection of the orbit; (b) $a_3\,, a_4$ projection of the same orbit. 88

30 The unstable manifold of p_C and its first iteration on the Poincaré section $a_1 = 0$. (a) a_2, a_3 projection of the unstable manifold on the Poincaré section; (b) One Poincaré iteration of the unstable manifold projected to a_2, a_3 components. 89

31 The unstable manifold of p_{R2} and its first iteration on the Poincaré section $a_4 = -0.122$. (a) a_2, a_3 projection of the unstable manifold on the Poincaré section; (b) One Poincaré iteration of the unstable manifold projected to a_2, a_3 components. 89

32 The return maps on the Poincaré sections using the intrinsic coordinates. (a) The return map for p_C on $a_1 = 0$; (b) The return map for p_{R2} on $a_4 = -0.122$. 90

33 An attracting cycle of the KSe near S_R with the period $T = 40.0565$ in the antisymmetric solution space with periodic boundary condition; $L = 38.5$ and $\nu = 1$. (a) a_1, a_2 Fourier modes projection of the UPO; (b) a_3, a_4 Fourier modes projection of the UPO. 90

34 A UPO of the KSe on S_R with the period $T = 60.1112$ in the antisymmetric solution space with periodic boundary condition; $L = 38.5$ and $\nu = 1$. (a) a_1, a_2 Fourier modes projection of the UPO; (b) a_3, a_4 Fourier modes projection of the UPO. 91

35 A non-symmetric UPO of the KSe on S_C with the period $T = 81.3345$ in the antisymmetric solution space with periodic boundary condition; $L = 38.5$ and $\nu = 1$. (a) a_1, a_2 Fourier modes projection of the UPO; (b) a_3, a_4 Fourier modes projection of the UPO. 91

36 A symmetric UPO of the KSe on S_R with the period $T = 77.4483$ in the antisymmetric solution space with periodic boundary condition; $L = 38.5$ and $\nu = 1$. (a) a_1, a_2 Fourier modes projection of the UPO; (b) a_3, a_4 Fourier modes projection of the UPO. 92

Abbreviations and Synonyms

SIS: Strange Invariant Set, an invariant set that has a fractal dimension.
UPO: Unstable Periodic Orbit.
cycle: a periodic orbit.
PDE: Partial Differential Equation.
ODE: Ordinary Differential Equation.
KSe: Kuramoto-Sivashinsky equation.
CGLe: Complex Ginzburg-Landau equation.
NSe: Navier-Stokes equation

SUMMARY

We propose a dynamical systems approach to study of turbulence (spatiotemporal chaos) based on the periodic orbit theory. The role of recurrent patterns and coherent structures is emphasized in the description. After a brief review of the periodic orbit theory in low dimensions, we discuss its possible application to the dynamics of spatially extended systems. The discussion is three-fold. First, we introduce a novel variational scheme for finding periodic orbits in general flows. Second, we check the periodic structures near the first instability of the complex Ginzburg-Landau equation and talk about its role in pattern formation. Third, we discuss the extensive numerical explorations that we have done to the Kuramoto-Sivashinsky system in the chaotic regime, including searching for periodic orbits, constructing symbolic dynamics and describing the chaotic invariant set in terms of intrinsic coordinates. In these investigations, we hope to identify and rank the nonlinear instabilities according to their relative importance.